



DEDICATED TO YOUR SMILE

Spodak Dental Group  
3911 W Atlantic Avenue  
Delray Beach, FL 33445  
561-498-0050

## Post-Operative Instructions Implant Surgery

### General Instructions:

- Avoid strenuous physical activity and smoking following the procedure.
- Start the prescribed medications immediately and avoid all alcoholic and carbonated beverages.
- The pain medications prescribed for you may cause drowsiness and impair your ability to drive and perform delicate tasks. Avoid all activities requiring your full, alert attention while on pain medication.
- Be sure to drink plenty of fluids during the healing process.
- Apply cold compresses to the surgical side of your face intermittently during the first 48 hours. Following this period, apply moist or wet warm compresses intermittently to reduce swelling as needed.
- If necessary, you can lubricate the corners of your mouth with lip balm.
- Begin a complete, but soft diet and avoid spicy and crunchy foods.

### Detailed Instructions:

- **Oral Hygiene:** Unless specifically instructed not to, on the day following surgery, light, gentle brushing of the teeth involved in the surgical area should be started. Please use a soft or extra soft tooth brush only. Frequent gentle rinsing of the mouth with warm saltwater (1 teaspoon salt with 8 oz. water) after the first 24 hours will aid in the healing process, and will help keep the mouth clean. If prescribed, utilize the bacterial mouth rinse (Peridex) twice daily and avoid eating or drinking for 20 minutes after each application. If given PerioSciences AO ProVantage Gel, apply five times a day for the first two weeks, and twice a day thereafter, preferably once in the morning and once in the evening.



DEDICATED TO YOUR SMILE

3911 West Atlantic Avenue | Delray Beach, FL 33445 [t] 561.498.0050 [f] 561.498.084 1

[www.SPODAKDENTAL.com](http://www.SPODAKDENTAL.com)

Dr. Daniel Torres  
[drtorres@spodakdental.com](mailto:drtorres@spodakdental.com)  
cell (954) 803-5221

- **Discomfort:** Following all types of surgery, a certain amount of discomfort is anticipated. If pain is not adequately controlled by the prescribed medication(s), or persists for a prolonged period of time (five days or more), please contact the office. Pain medication is prescribed to alleviate discomfort during the initial healing period. However, patients may elect to not take this medication at all, reduce the frequency or dosage, or substitute the prescribed pain medication with over the counter medications, such as Tylenol, Ibuprofen (Advil, Motrin), or Aleve.
- **Swelling:** A slight amount of swelling and discomfort is not unusual following most oral surgery and usually peaks two days after surgery. The use of cold compresses gently applied to the surgical area for five to 10 minutes every 30 minutes (during the first 24-48 hours) will help to control the post-operative swelling and discomfort. If swelling persists, apply moist heat (warm washcloth or hot water bottle). If swelling continues after two days, or interferes with swallowing, please call the office.
- **Bleeding:** Some bleeding or oozing may stain your saliva. If the bleeding is excessive, (i.e. nose bleed), for more than five minutes, two hours after your procedure, continue to bite on gauze. Avoid exertion, lie down with your head elevated at least 30 degrees (two pillows), and call the office.
- **Fever:** A slight fever or chills may occur. It is of no consequence unless it persists more than 24 hours. A persistent low grade temperature or one above 102 degrees should be reported to our office.
- **Diet:** Drink fluids immediately after surgery and avoid chewing on the operated area of your mouth. Soft foods can be eaten until you feel ready to return to your regular diet. The following require little pressure to chew and provide the necessary nutrients: protein shakes, bananas, cooked fish, cereals, pasta, mashed potatoes and whole grain soft breads. You should continue your usual vitamin supplements. Returning to your normal diet and proper calorie intake will promote faster healing.

### Special Medications:

- If you are taking other medication prescribed by your physician, be sure to inform your doctor and continue taking them as usual unless otherwise directed. Antibiotics or other medication(s) may be prescribed for you and please take these as directed on the label(s). Antibiotics especially need to be taken as prescribed and one should complete the full regimen as given. Female patients should also be aware that some antibiotics may interfere with the effectiveness of certain oral contraceptives.



DEDICATED TO YOUR SMILE

3911 West Atlantic Avenue | Delray Beach, FL 33445 [t] 561.498.0050 [f] 561.498.084 1

Dr. Daniel Torres  
 drtorres@spodakdental.com  
 cell (954) 803-5221

[www.SPODAKDENTAL.com](http://www.SPODAKDENTAL.com)

### **Specific Post Implant Placement Instructions:**

- Following your surgical procedure, extreme care should be exercised to avoid placing any biting pressure on the implant or implant site(s), no matter how light the biting force. If a removable prosthetic appliance was placed, it should not contact the implant or tissue in the area of the implant placement at all. If a temporary was placed on the implant, all biting contact with or without food should be avoided. If you feel any contact on the temporary when your teeth come completely together, please contact the office immediately and schedule an appointment right away. Premature pressure on an implant prior to three months of healing may result in implant failure and/or healing complications.
- Remember that not all implants fuse to the surrounding bone, however, following all the instructions will help minimize the risk of implant failure.

### **Future Appointments:**

- Appointments will be made for you to evaluate your healing and monitor your progress following your surgical procedure(s). These appointments are important and should be kept as scheduled.



DEDICATED TO YOUR SMILE

3911 West Atlantic Avenue | Delray Beach, FL 33445 [t] 561.498.0050 [f] 561.498.0841

Dr. Daniel Torres  
drtorres@spodakdental.com  
cell (954) 803-5221

[www.SPODAKDENTAL.com](http://www.SPODAKDENTAL.com)